

Prophylactic Antibiotic Regimen Selection for Surgery

Surgical Procedure	Approved Antibiotics
CABG, Other Cardiac or Vascular	Cefazolin, Cefuroxime or Vancomycin** If β -lactam allergy: Vancomycin* or Clindamycin*
Hip/Knee Arthroplasty	Cefazolin or Cefuroxime or Vancomycin** If β -lactam allergy: Vancomycin* or Clindamycin*
Colon	Cefotetan, Cefoxitin, or Ampicillin/Sulbactam or Ertapenem† OR Cefazolin or Cefuroxime + Metronidazole If β -lactam allergy: Clindamycin + Aminoglycoside, or Clindamycin + Quinolone, or Clindamycin + Aztreonam OR Metronidazole with Aminoglycoside, or Metronidazole + Quinolone
Hysterectomy	Cefotetan, Cefazolin, Cefoxitin, Cefuroxime, or Ampicillin/Sulbactam If β -lactam allergy: Clindamycin + Aminoglycoside, or Clindamycin + Quinolone, or Clindamycin + Aztreonam OR Metronidazole with Aminoglycoside, or Metronidazole + Quinolone

Prophylactic Antibiotic Regimen Selection for Surgery

Surgical Procedure	Approved Antibiotics
Special Considerations	<p>*For cardiac, orthopedic, and vascular surgery, if the patient is allergic to β-lactam antibiotics, Vancomycin or Clindamycin are acceptable substitutes.</p> <p>**Vancomycin is acceptable with a physician/ APN/PA/pharmacist documented justification for its use (see data element Vancomycin)</p> <p>† A single dose of ertapenem is recommended for colon procedures.</p>

The antibiotic regimens described in this table reflect the combined, published recommendations of the Specifications Manual for discharges from 10-01-09 (4Q09) through 03-31-10 (1Q10)



This material provided by Quality Insights of Delaware, the Medicare Quality Improvement Organization for Delaware, was prepared by the Oklahoma Foundation for Medical Quality, under contract with the Centers for Medicare & Medicaid Services (CMS), an agency of the U.S. Department of Health and Human Services. The contents presented do not necessarily reflect CMS policy. Publication number 9SOW-DE-PASA-09.80. Rev. 9/09.